



MIXED FERTILIZER PROTOCOL FOR RICE [TRANSPLANTING METHOD]

GROWTH STAGE	SEEDING	SEEDLING	TILLERING		REPRODUCTIVE	HEADING
						
DAT - Days After Transplanting	3 Days	4-8 DAT	12-16 DAT	20-24 DAT	50-54 DAT	62-66 DAT
INTERVAL (Between Applications)	Before Pulling	8 to 12 days	8 to 10 days	8 to 10 days	One month	12 to 14 days
DOSAGE PER 16L TANKLOAD (The XANADU bottle cap holds approximately 25ml up to the topmost screw line)  = 25 ml						
XANADU F1 MAXPOWER	200 ml	BASAL RECOMMENDATION: UREA (46-0-0) 50 Kg + T14 (14-14-14) 50 Kg	200 ml			
XANADU F2 MAXPOWER				200 ml		
				ADD DURING DRY SEASON:		
				 +100 ml		
XANADU F3 MAXPOWER					250 ml	250 ml
NO. OF TANKLOADS	1 / Seedbed		9/ Hectare	10/ Hectare	10/ Hectare	10/ Hectare
DOSAGE PER HECTARE						
XANADU MAXPOWER	200 ml F1		1.8 L F1	2 L F2	2.5L F3	2.5L F3
				ADD DURING DRY SEASON:		
				 +1L F2		
BASE REQUIREMENT / HA: 2L F1 + 2L F2 + 5L F3						



REMINDER:

Follow the recommended increase in dosage if the weather is dry and the sunlight intensity is high at 20-24 DAT.